



Shin, M. J., Cho, Y., & Davey Smith, G. (2017). Alcohol consumption, aldehyde dehydrogenase 2 gene polymorphisms, and cardiovascular health in Korea. *Yonsei Medical Journal*, 58(4), 689-696.  
<https://doi.org/10.3349/ymj.2017.58.4.689>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY-NC

Link to published version (if available):  
[10.3349/ymj.2017.58.4.689](https://doi.org/10.3349/ymj.2017.58.4.689)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



## Alcohol Consumption, *Aldehyde Dehydrogenase 2* Gene Polymorphisms, and Cardiovascular Health in Korea

Min-Jeong Shin<sup>1</sup>, Yoonsu Cho<sup>1,2</sup>, and George Davey Smith<sup>2</sup>

<sup>1</sup>Department of Public Health Sciences, BK21PLUS Program in Embodiment: Health-Society Interaction, Graduate School, Korea University, Seoul, Korea;

<sup>2</sup>MRC Integrative Epidemiology Unit, School of Social & Community Medicine, University of Bristol, Bristol, UK.

Alcohol consumption is a serious health issue in Korea in terms of the amount consumed and the behavior related to its consumption. Aldehyde dehydrogenase 2 (ALDH2) is a key enzyme in alcohol metabolism that degrades acetaldehyde to nontoxic acetic acid. The enzyme is coded by the *ALDH2* gene, which is commonly polymorphic in East Asian populations. A point mutation in the *ALDH2* gene (the rs671 allele) yields an inactive form of ALDH2 that causes acetaldehyde accumulation in the body after alcohol consumption, thereby inhibiting normal alcohol metabolism. Individuals who are homozygous for polymorphism in *ALDH2* tend to refrain from drinking alcohol, decreasing their chances of developing alcoholism and exposure to the associated risks. Mendelian randomization (MR) studies have demonstrated that alcohol consumption predicted by *ALDH2* genotype is causally related to cardiovascular risks. Moreover, recent MR studies suggest that the *ALDH2* variant has mechanistic effects on some disease outcomes or mortality through increased blood levels of acetaldehyde, showing differences therein between heterozygotes (*ALDH2*\*2\*2) and homozygotes (*ALDH2*\*1\*2) in those who consume alcohol. Accordingly, consideration of *ALDH2* genotype in alcohol prevention programs is warranted. In conclusion, strategies that incorporate genetic information and provide an evidential basis from which to help people make informed decisions on alcohol consumption are urgently required.

**Key Words:** Alcohol, aldehyde dehydrogenase 2, Mendelian randomization analysis, guidelines

### INTRODUCTION

In Korea, alcohol abuse is a serious health issue in terms of the amount consumed and behaviors related to its consumption.<sup>1</sup> According to the Korea National Health and Nutrition Examination Survey (KNHANES) 2013,<sup>2</sup> the percentage of individuals that consumed alcohol at least once a month comprised approximately 55% in 2005 and 60% in 2013, showing an increasing trend. Heavy drinkers (those who consume at least three glasses of alcohol per day) reportedly accounted

for one in four men, while men in their 40s and 50s exhibited the highest alcohol abuse and dependency rates. In addition, the KNHANES revealed that the high-risk drinking rates by sex, as well as the proportion of youths exposed to alcohol, have gradually increased.<sup>2</sup>

It is generally accepted that alcohol abuse has negative effects on health and quality of life; thus, guidelines for the regulation of alcohol consumption at a global level have been suggested. An example of this would be the “Global Strategy to Reduce the Harmful Use of Alcohol” proposed by the World Health Organization.<sup>3</sup> In the United States, alcohol prevention programs and services, such as AlcoholEdu and Project ALERT, have been initiated to regulate alcohol-associated risks.<sup>4</sup> In Korea, however, although an age limit for alcohol consumption is in effect, there are no specific regulations on the time and place of alcohol sales. Moreover, South Korea has no national agency or independent research institution for developing guidelines and policies on alcohol consumption. The absence of a detailed national policy on alcohol consumption in Korea has led to a culture of binge drinking and an increase in the

**Received:** March 10, 2017 **Revised:** April 27, 2017

**Accepted:** May 2, 2017

**Corresponding author:** Dr. Min-Jeong Shin, Department of Public Health Sciences, Graduate School, Korea University, 145 Anam-ro, Seongbuk-gu, Seoul 02841, Korea. Tel: 82-2-3290-5643, Fax: 82-2-940-2849, E-mail: mjshin@korea.ac.kr

•The authors have no financial conflicts of interest.

© Copyright: Yonsei University College of Medicine 2017

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

incidence of alcohol-related accidents. Therefore, we find it necessary to focus on the control of alcohol abuse in this population.

It has been suggested that individual alcohol sensitivity and its influence are affected by genotype. The *aldehyde dehydrogenase (ALDH2)* gene encodes the major enzyme responsible for alcohol metabolism, which eliminates acetaldehyde.<sup>5</sup> The *ALDH2* gene is polymorphic in East Asians, including Koreans, and influences blood levels of acetaldehyde and behavior after consumption, which is associated with the risk for alcoholic cirrhosis, cardiovascular disease (CVD), and cancer.<sup>6-10</sup> The *ALDH2* polymorphism is associated with a flushing response to alcohol use and lowers alcohol consumption, which acts as protective factors for CVDs.<sup>6,11</sup> A previous study has reported that *ALDH2\*2* homozygotes and heterozygotes had 18 and 5 times higher blood alcohol levels, respectively, than *ALDH2\*1* homozygotes after drinking the same amount of alcohol.<sup>12</sup> Thus, it has been suggested that intervention for alcohol use should be individually specified, considering genotype information. In this review, we highlight the role of *ALDH2* polymorphisms in alcohol-related health problems in Koreans. We also propose an educational model incorporating genetic information on *ALDH2* that may improve alcohol consumption-related behaviors in this population.

## ALCOHOL: A MAJOR PREVENTABLE RISK FOR HEALTH PROBLEMS

A number of epidemiologic studies have found that alcohol intake has both positive and negative effects on cardiovascular health. It has been generally reported that high alcohol intake is a risk factor for diabetes and CVDs.<sup>13,14</sup> On the other hand, other epidemiological evidence have suggested that moderate alcohol consumption might have a protective effect against chronic diseases, reflected in the so-called J-shaped association.<sup>15</sup> However, the results of observational studies have limitations because they are affected by confounding variables or reverse causation when investigating the association between alcohol consumption and the risk for chronic diseases.<sup>16</sup> For example, people who stopped drinking, used as a control for moderate drinkers, may do so because of health problems, indicating reverse causation. In supporting this, Ng Fat, et al.<sup>17</sup> confirmed that poor health directly influenced nondrinking throughout the life course, indicating a sick-quitter bias. Consistently, a recent meta-analysis<sup>18</sup> found that a reduction in mortality risk in moderate drinkers (one to two glasses a day) was not supported when they were compared with intermittent drinkers (at most one glass a week/intake of less than 1.3 g per day) or those who never drank. In other words, the lower mortality risk in intermittent drinkers can be explained by reverse causation or confounding variables, rather than by the amount of alcohol intake. Hence, it is difficult to conclude that lowering

alcohol intake reduces mortality. Furthermore, it has also been reported that there is no association between limited alcohol intake and cancer- or cardiovascular-related mortality<sup>19</sup> and that even moderate drinking can be harmful.<sup>20</sup> For these reasons, the International Agency for Research on Cancer classified alcohol as a Group 1 carcinogen,<sup>21</sup> while the European Union enhanced their cancer prevention guidelines to include “no alcohol drinking” in 2014.<sup>22</sup> Following these global recommendations, the Ministry of Health and Welfare of South Korea revised their guidelines in March 2016 to “To prevent cancer, drinking even small amounts of alcohol (one or two glasses a day) should be avoided,” which was previously “Drinking up to two glasses a day.”<sup>23</sup> The results of a recent meta-analysis in South Korea also showed that moderate alcohol intake has no beneficial effect on total mortality,<sup>19</sup> supporting current considerations to extend the application of this drinking guideline to other chronic diseases, such as stroke. However, data are scarce on the relationship between alcohol consumption and chronic diseases, such as diabetes and CVD, in Korea, and information on safe levels of alcohol consumption for diseases of interest is hard to find. Thus, it is necessary to accumulate scientific evidence on the effects of alcohol consumption on health outcomes in order to establish a health policy that promotes a desirable alcohol intake and drinking pattern in this population.

## THE EPIDEMIOLOGY OF GENETIC DIFFERENCES IN ALCOHOL USE: *ALDH2* AS A GENETIC MARKER FOR ALCOHOL CONSUMPTION


Alcohol consumption behavior is considered to reflect complex etiology, involving environmental (e.g., cultural, social, and political) and genetic factors.<sup>24-26</sup> Although some studies of alcohol use support this general understanding,<sup>27,28</sup> there is evidence that genetic variations, such as *ALDH2\*2* and *alcohol dehydrogenase 1B (ADH1B)\*2*, have measurable effects on alcohol consumption in East Asians.<sup>29-31</sup> Consistently, a genome-wide association study in a Korean population identified genetic loci associated with alcohol intake.<sup>32</sup> According to this report, genes, including *C12orf51*, *CCDC63*, *MYL2*, *OAS3*, *CUX2*, and *RPH3A*, are in high linkage disequilibrium (LD) with the *ALDH2* gene and are associated with amounts of alcohol consumed. Particularly, a genetic polymorphism in *C12orf51* (rs2074356) at 12q24.13, which was in substantial LD with several intronic single nucleotide polymorphisms in *ALDH2*, such as rs2238151, showed the highest correlation with alcohol consumption and was associated with delayed acetaldehyde catabolism.<sup>33</sup> Another study in a Korean population reported that *ADH1B\*47Arg* and *ALDH2\*487Glu*, which affect alcohol metabolism, explained 86.5% of Korean alcoholics.<sup>34</sup> This study also suggested that individuals who have

the *ADH1B*\*47Arg allele and the susceptible *ALDH2*\*487Glu allele at either loci had 11 times greater risk for alcoholism. Based on these observations, it is possible that these two genes, which are closely related to alcohol metabolism, could well explain the reason for alcohol intake and the probability of becoming an alcoholic in Koreans. Several genetic polymorphisms in *ADH1B* and *ALDH2* reportedly generate functional differences in the activities of enzymes involved in alcohol metabolism, which influence the level of acetaldehyde, an intermediate metabolite of alcohol in the body. Generally, *ALDH2* variation has a strong effect on alcohol metabolism, affecting blood levels of acetaldehyde to a greater extent than *ADH1B* (Fig. 1A)<sup>29,35,36</sup> A study in a Korean population reported the lack of a significant association between the *ADH1B* genotype and self-rating of the effects of alcohol.<sup>36</sup> In line with this, a previous study in Asian-Americans also showed that the *ALDH2*\*2 variant had a protective effect on alcohol use behavior, whereas the *ADH1B* variant was not associated with alcohol consumption.<sup>35</sup> However, this requires further confirmation in other populations. Additionally, in East Asian populations (Chinese, Japanese, Korean, and Thai), the possession of two *ALDH2*\*2 alleles was associated with a nine-fold reduction in alcohol consumption, whereas the possession of two *ADH1B*\*2 alleles was associated with a five-fold reduction.<sup>29</sup> Thus, particular attention has been focused on the roles of genetic variants in *ALDH2* in alcohol metabolism. Interestingly, reports show that up to 50% of East Asians<sup>9,37,38</sup> and about 25–35% of Koreans have an inactive form of *ALDH2*,<sup>39,40</sup> which is scarce in North Americans or Europeans (Fig. 1B). The inactive form of *ALDH2* due to genetic variation causes the accumulation of acetaldehyde in the body, which leads to physiologic adverse effects, such as flushing,


palpitations, headache, and displeasure,<sup>41</sup> as well as a behavioral influence in refraining from alcohol consumption. This genetically chosen behavior helps lower the chances of being alcoholic and reduces exposure to its associated risks. In this regard, several studies have proposed the use of *ALDH2* as a biomarker for alcoholism.<sup>41,42</sup>

## ACETALDEHYDE METABOLISM, PHYSIOLOGICAL RESPONSES, AND HEALTH EFFECTS OF *ALDH2* VIA ALCOHOL CONSUMPTION ACCORDING TO *ALDH2* VARIANTS

Numerous studies have reported differences in alcohol metabolism, physiological functions, and drinking behavior according to the rs671 genetic polymorphism. Specifically, heterozygous individuals (*ALDH2*\*1/\*2) exhibit slower acetaldehyde degradation rates, equating to markedly higher levels of acetaldehyde than in homozygotes (*ALDH2*\*1/\*1),<sup>5</sup> which was also verified in Koreans.<sup>43,44</sup> It is generally accepted that high acetaldehyde levels in the blood exert negative effects on cardiovascular functions, causing subsequent elevations in heart rate, cardiac output, and blood flow velocity in the facial artery.<sup>45</sup> Also, the accumulation of acetaldehyde over time induces various symptoms, including facial flushing.<sup>46</sup> According to a survey of 250 Thai people,<sup>47</sup> 80% of those with the *ALDH2*\*1/\*2 genotype responded that they experienced flushing immediately after drinking alcohol. In contrast, only 28% of those with the *ALDH2*\*1/\*1 genotype answered that they experienced flushing immediately after alcohol consumption. This was further supported by a clinical study<sup>48</sup> showing that the *ALDH2*\*1/\*2 genotype raised the heart rate and induced facial flushing in



**Fig. 1.** Alcohol metabolism according to genetic polymorphisms. (A) Aldehyde dehydrogenase 2 (*ALDH2*) and alcohol dehydrogenase 1B (*ADH1B*) are involved in alcohol metabolism. Generally, *ALDH2* variants have a greater influence on alcohol intake than *ADH1B* variants in East Asians. (B) *ALDH2* variants mainly exist in East Asian populations. Adapted from Cho, et al. Sci Rep 2015;5:18422.<sup>6</sup>



**Fig. 2.** Scheme of the Mendelian randomization approach. The scheme of the Mendelian randomization approach. (A) This method uses a genetic variant, which proxies for exposure that is expected to be related to the outcome. (B) Using the Mendelian randomization approach, the causal effects of alcohol intake on cardiovascular outcomes can be investigated while controlling confounders and reverse causation. *ALDH2*, aldehyde dehydrogenase 2.

individuals after alcohol consumption relative to those with the *ALDH2\*1/\*1* genotype. Moreover, there is also evidence that the *ALDH2\*1/\*2* genotype aggravates psychomotor functions relative to the *ALDH2\*1/\*1* genotype in Koreans.<sup>49,50</sup>

While people with the inactive form of ALDH2 tend to refrain from alcohol consumption, thus decreasing their exposure to associated risks, the inactive form of ALDH2 also manifests negative effects on health because of the toxicity of accumulated acetaldehyde after drinking. So far, studies on an ALDH2 variant (the rs671 allele) have suggested that an alteration in ALDH2 activity caused by genetic variation is associated with a 20–60% increase in susceptibility to CVD and cancer in Asians.<sup>51–54</sup> However, previous studies have not considered the possible association between alcohol drinking behavior caused by the ALDH2 variant and CVD.<sup>55</sup> Similarly, associations observed by Gu and Li<sup>52</sup> and Wang, et al.<sup>53</sup> between the *ALDH2\*504lys* allele and increased risk of myocardial infarction were based on ordinary least square estimates. In addition, an association study in a Chinese population reported that the *ALDH2 Glu504Lys* polymorphism was associated with the risk for hypertension in men, but not in women, who consumed relatively lower amounts of alcohol independent of their genotype.<sup>54</sup> Such a difference in the association between men and women provides evidence that the *ALDH2* polymorphism influences outcomes only through alcohol intake.<sup>56</sup> This is in line with recent studies showing that lower alcohol intake in individuals with the rs671 A allele, compared to wild-type homozygotes,<sup>57,58</sup> was associated with reduced cardiometabolic diseases, glucose levels, and blood pressure in men, who generally consumed more alcohol than women.

It was recently reported that Koreans with the inactive form of ALDH2 had a higher incidence of myocardial infarction due to the accumulation of acetaldehyde.<sup>59</sup> However, alcohol intake had not been taken into account in the evaluation of the association between ALDH2 and CVD. Additionally, another study in the Korean population reported a 52% lower incidence of lung cancer in nondrinkers with the inactive form of ALDH2 than those without.<sup>60</sup> Based on evidence that *ALDH2\*2\*2* homozygotes consume considerably less alcohol than the wild-type homozygotes (*ALDH2\*1*), with heterozygotes consuming intermediate amounts,<sup>8</sup> it might be more reasonable to consider *ALDH2* as a proxy for alcohol intake when investigating the effect of *ALDH2* variants on CVD outcomes.

## MENDELIAN RANDOMIZATION ANALYSIS BASED ON *ALDH2* POLYMORPHISMS

Recently, our research group tested whether alcohol intake is associated with cardiovascular risks in the Korean population, using genetic variation in *ALDH2* (rs671) as a proxy for alcohol intake, in a MR framework (Fig. 2).<sup>6</sup> We demonstrated, for

the first time in a Korean population, a causal relationship between alcohol intake and hypertension and several cardiometabolic risk factors using an MR method that eliminated interference by confounders.<sup>61</sup> Many epidemiologic studies have shown that moderate alcohol intake protects individuals against CVDs, as reflected by the J-shaped correlation,<sup>15</sup> are limited by confounding variables, including smoking and socioeconomic position, and reverse causation due to the characteristics of observational studies. For example, light or moderate alcohol users also tend to have other characteristics that are particularly beneficial to health, such as high income levels, good education, and regular exercise (confounding). Also, people diagnosed with chronic diseases tend to change their alcohol consumption patterns over time (reverse causation). In our previous study utilizing the MR framework,<sup>6</sup> we found that the amount and pattern of alcohol intake are influenced by genotype and are associated with an increased risk for CVDs in a dose-dependent manner throughout life. Our findings are consistent with those of a study using the MR method in Chinese individuals,<sup>11</sup> which showed that alcohol intake is predicted by genetic polymorphisms in *ALDH2* and has negative effects on blood pressure and blood triglyceride levels. It has yet to be determined whether the effects of genetic polymorphisms in *ALDH2* on risk factors for CVD shown in our study, as well as those in other populations, are caused by the physiologic effects of these genetic variations or a subsequent result of alcohol intake (Table 1). A recent study in South Korea showed that genetic polymorphisms in *ALDH2* are correlated with stroke in men only,<sup>62</sup> raising the possibility that the observed association is mainly attributable to the strong correlation mediated by alcohol intake, especially in men. Further, very recently, it has been suggested that alcohol flushing can be used as an instrumental variable to evaluate the health impact of alcohol consumption, since alcohol flushing is closely related to ALDH2 deficiency.<sup>63</sup> Using a phenotypic proxy of a genotype allows researchers to achieve MR-inspired causal inference even genetic data are absent.<sup>64</sup>

## A GENETIC EDUCATION MODEL FOR ALCOHOL CONSUMPTION AND RELATED BEHAVIOR

According to a recent report in South Korea, the mortality rate of Korean men with chronic diseases, such as oral or laryngeal cancer, is approximately 30–40% and is attributable to alcohol consumption.<sup>65</sup> In order to reduce alcohol intake more effectively at the population level, novel personalized drinking guidelines incorporating genetic information, such as *ALDH2* variants, would be of value. It was previously reported that *ALDH2* variants increased the risk of head and neck cancer only with the interaction with drinking alcohol, and *ALDH2\*1\*2* heterozygotes showed a less severe reaction to alcohol but an



**Table 1.** Mendelian Randomization Studies on Alcohol Consumption Predicted by Genotype and CVD Risk Factors

Author	Population	Instrument	Outcome	Statistics	Major results
Jee, et al. <sup>69</sup>	Korean (n=2993 men and 1374 women)	rs671 in <i>ALDH2</i>	Serum glucose levels	Instrumental variable regression analysis	An increase of 1 unit of alcohol with the G allele of rs671 increased serum glucose levels in men.
Cho, et al. <sup>6</sup>	Korean (n=3365 men and 3787 women)	rs671 in <i>ALDH2</i>	Risk factors for CVD	Instrumental variable regression analysis	An increase of 1 unit of alcohol with the G allele of rs671 increased CVD risk factors, including blood pressure, waist-to-hip ratio, fasting blood glucose, and TG levels.
Xu, et al. <sup>70</sup>	Chinese (n=2321 men and 2757 women)	rs671 in <i>ALDH2</i>	Gamma- glutamyltransferase	Instrumental variable regression analysis	An increase of 1 unit of alcohol with the G allele of rs671 increased gamma-glutamyltransferase by 10.60 U/L per 10 g ethanol/day.
Au Yeung, et al. <sup>71</sup>	Chinese (n=4588 men)	rs671 in <i>ALDH2</i>	Heart rate	Instrumental variable regression analysis	An increase of 1 unit of alcohol with active <i>ALDH2</i> was associated with an increase in heart rate of 0.98 beats per minute.
Au Yeung, et al. <sup>72</sup>	Chinese (n=4867 men)	rs671 in <i>ALDH2</i>	Self-reported CVD and biologic CVD risk factors	Instrumental variable regression analysis	An increase of 1 unit of alcohol with the G allele of rs671 was associated with an increase in HDL cholesterol per alcohol unit, and DBP, LDL cholesterol, TG, or glucose, and the prevalence of self-reported CVD.
Au Yeung, et al. <sup>73</sup>	Chinese (n=4707 men for alcohol consumption was not associated with delayed 10-word recall score; 2284 men for MMSE score)	rs671 in <i>ALDH2</i>	Cognitive function	Instrumental variable regression analysis	An increase of 1 unit of alcohol with the G allele of rs671 was not associated with delayed 10-word recall score or MMSE score.
Tabara, et al. <sup>74</sup>	Japanese (n=2756 men and 5608 women)	<i>ALDH2</i>	LDL/HDL cholesterol levels	Linear regression analysis with an additive genetic model	The <i>ALDH2</i> *1 variant was strongly associated with higher alcohol consumption in men. The <i>ALDH2</i> *1 appeared causally related to increased HDL cholesterol levels in men. The <i>ALDH2</i> variant also related to decreased total LDL cholesterol levels, while the LDL profiles were worsened.
Tabara, et al. <sup>75</sup>	Japanese (n=2289 men and 1940 women)	<i>ALDH2</i>	LDL/HDL cholesterol levels	Linear regression analysis with an additive genetic model	The <i>ALDH2</i> *1 variant was strongly associated with higher alcohol consumption in men. The <i>ALDH2</i> *1 appeared causally related to increased HDL cholesterol levels and reduced LDL cholesterol levels in men. Alcohol consumption was related to not only increased HDL cholesterol levels but also decreased LDL cholesterol levels and particle numbers.
Taylor, et al. <sup>11</sup>	Chinese (n=1712 diabetes cases and 2076 controls)	rs671 in <i>ALDH2</i>	Cardiovascular and metabolic factors	Linear regression analysis with an additive genetic model	The A allele of rs671 was strongly associated with reduced alcohol use in men and women. The A allele was associated with lower blood pressure, lower HDL cholesterol and lower triglycerides in men.

*ALDH2*, aldehyde dehydrogenase 2; CVD, cardiovascular disease; DBP, diastolic blood pressure; LDL, low-density lipid; HDL, high-density lipid; TG, triglyceride; MMSE, Mini-Mental State Examination.

The table only includes Mendelian randomization studies based on instrumental variable (IV) regression using rs671 in *ALDH2* as an instrument. Papers included in Table 1 were collected from MEDLINE (<http://www.ncbi.nlm.nih.gov/PubMed>) using the following keywords: "Aldehyde dehydrogenase 2" and "*ALDH2*" in combination with "Mendelian randomization" and "alcohol." The abstracts of the collected articles were scanned for inclusion of any cardiovascular-related outcomes of interest. We searched for all papers published before January 2017. The minimum requirement for inclusion was Mendelian randomization analysis using *ALDH2* as an instrument. We excluded meta-analyses, because they did not perform IV regression analysis directly. Among the 20 papers identified by the search strategy, irrelevant articles (n=7) and meta-analyses (n=4) were excluded. When sufficient information could not be obtained, the study was excluded.

elevated risk for cancer with the same amount of alcohol.<sup>8</sup> In support of this, an animal study demonstrated that maternal deficiency in ALDH2 due to the *ALDH2\*2* allele resulted in extensive developmental malformations, particularly in cases of alcohol consumption during pregnancy. They also reported that genotoxic aldehydes in *ALDH2\*2* homozygous fetuses born from *ALDH2\*2* homozygous or heterozygous mothers who consumed alcohol were continuously accumulated, possibly resulting in birth defect.<sup>66</sup> Given that approximately 30% of the Korean population are genetically deficient in *ALDH2*,<sup>6</sup> it would be effective if people could be informed about their own genetic predisposition to enable them to identify their own cardiovascular risks. A previous study on Health Risk Appraisal Models, which made use of information on *ALDH2* variants to evaluate disease outcomes, supported this notion, showing high detection rate.<sup>67</sup> In line with this, the inclusion of genetic information is now regarded as a very useful tool for educating alcohol intake and drinking behavior properly in community and/or clinical settings,<sup>68</sup> yet still, such a genetic education model is not available in Korea.

## CONCLUSION

Although about 30% of Koreans have the inactive form of ALDH2, alcohol intake rates are still high in Koreans. This genetically high-risk group represents a target for a customized alcohol-prevention program. In particular, the development of strategies using genetic information may prove necessary to modifying alcohol consumption behaviors in Korea. This novel approach is expected allow people to make informed decision about their own alcohol consumption and, ultimately, raise public awareness of alcohol-gene interactions that is associated with harmful drinking behavior and health outcomes.

## ACKNOWLEDGEMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2015R1A2A1A15054758 to MJS).

YC, and GDS work within the Medical Research Council Integrative Epidemiology Unit at the University of Bristol (MC\_UU\_12013/1).

## REFERENCES

1. Korea Centers for Disease Control and Prevention. Development of surveillance system on alcohol consumption and alcohol-related harms based on community health survey. Cheongju: Korea Centers for Disease Control and Prevention; 2011.
2. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. Korea Health Statistics 2013: Korea National Health and Nutrition Examination Survey (KNHANES VI-1). Cheongju: Korea Centers for Disease Control and Prevention; 2014.
3. World Health Organization. Global strategy to reduce the harmful use of alcohol. Geneva: World Health Organization; 2010.
4. Foxcroft DR, Lister-Sharp D, Lowe G. Alcohol misuse prevention for young people: a systematic review reveals methodological concerns and lack of reliable evidence of effectiveness. *Addiction* 1997;92:531-7.
5. Chen YC, Peng GS, Wang MF, Tsao TP, Yin SJ. Polymorphism of ethanol-metabolism genes and alcoholism: correlation of allelic variations with the pharmacokinetic and pharmacodynamic consequences. *Chem Biol Interact* 2009;178:2-7.
6. Cho Y, Shin SY, Won S, Relton CL, Davey Smith G, Shin MJ. Alcohol intake and cardiovascular risk factors: a Mendelian randomisation study. *Sci Rep* 2015;5:18422.
7. Boccia S, Hashibe M, Galli P, De Feo E, Asakage T, Hashimoto T, et al. Aldehyde dehydrogenase 2 and head and neck cancer: a meta-analysis implementing a Mendelian randomization approach. *Cancer Epidemiol Biomarkers Prev* 2009;18:248-54.
8. Lewis SJ, Smith GD. Alcohol, ALDH2, and esophageal cancer: a meta-analysis which illustrates the potentials and limitations of a Mendelian randomization approach. *Cancer Epidemiol Biomarkers Prev* 2005;14:1967-71.
9. Yokoyama A, Omori T, Yokoyama T. Alcohol and aldehyde dehydrogenase polymorphisms and a new strategy for prevention and screening for cancer in the upper aerodigestive tract in East Asians. *Keio J Med* 2010;59:115-30.
10. He L, Deng T, Luo H. Aldehyde dehydrogenase 2 (ALDH2) polymorphism and the risk of alcoholic liver cirrhosis among East Asians: a meta-analysis. *Yonsei Med J* 2016;57:879-84.
11. Taylor AE, Lu F, Carslake D, Hu Z, Qian Y, Liu S, et al. Exploring causal associations of alcohol with cardiovascular and metabolic risk factors in a Chinese population using Mendelian randomization analysis. *Sci Rep* 2015;5:14005.
12. Enomoto N, Takase S, Yasuhara M, Takada A. Acetaldehyde metabolism in different aldehyde dehydrogenase-2 genotypes. *Alcohol Clin Exp Res* 1991;15:141-4.
13. Matsumoto C, Miedema MD, Ofman P, Gaziano JM, Sesso HD. An expanding knowledge of the mechanisms and effects of alcohol consumption on cardiovascular disease. *J Cardiopulm Rehabil Prev* 2014;34:159-71.
14. Knott C, Bell S, Britton A. Alcohol consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of more than 1.9 million individuals from 38 observational studies. *Diabetes Care* 2015;38:1804-12.
15. Corrao G, Rubbiati L, Bagnardi V, Zambon A, Poikolainen K. Alcohol and coronary heart disease: a meta-analysis. *Addiction* 2000;95:1505-23.
16. Freiberg MS, Samet JH. Alcohol and coronary heart disease: the answer awaits a randomized controlled trial. *Circulation* 2005;112:1379-81.
17. Ng Fat L, Cable N, Marmot MG, Shelton N. Persistent long-standing illness and non-drinking over time, implications for the use of lifetime abstainers as a control group. *J Epidemiol Community Health* 2014;68:71-7.
18. Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ* 2011;342:d671.
19. Park JE, Choi TY, Ryu Y, Cho SI. The relationship between mild alcohol consumption and mortality in Koreans: a systematic review and meta-analysis. *BMC Public Health* 2015;15:918.
20. Holmes MV, Dale CE, Zuccolo L, Silverwood RJ, Guo Y, Ye Z, et al. Association between alcohol and cardiovascular disease: Mendelian randomization study. *BMJ* 2014;349:g5000.

- lian randomisation analysis based on individual participant data. *BMJ* 2014;349:g4164.
21. International Agency for Research on Cancer. Agents classified by the IARC Monographs, volumes 1-104 [accessed on 2017 January 5]. Available at: <http://monographs.iarc.fr/ENG/Classification/ClassificationsAlphaOrder.pdf>.
  22. Scoccianti C, Cecchini M, Anderson AS, Berrino F, Boutron-Ruault MC, Espina C, et al. European code against cancer 4th edition: alcohol drinking and cancer. *Cancer Epidemiol* 2015;39 Suppl 1: S67-74.
  23. Korea Centers for Disease Control and Prevention, National Cancer Center. Guideline for prevention of cancer (Korean). Cheongju: Korea Centers for Disease Control and Prevention; 2016.
  24. Buscemi L, Turchi C. An overview of the genetic susceptibility to alcoholism. *Med Sci Law* 2011;51 Suppl 1:S2-6.
  25. Dragan WL, Oniszczenko W, Czerski PM, Dmitrzak-Weglarczyk M. [Family-based association study of dopaminergic genes polymorphisms and EAS temperamental traits]. *Psychiatr Pol* 2013;47: 185-95.
  26. Heath AC, Bucholz KK, Madden PA, Dinwiddie SH, Slutske WS, Bierut LJ, et al. Genetic and environmental contributions to alcohol dependence risk in a national twin sample: consistency of findings in women and men. *Psychol Med* 1997;27:1381-96.
  27. Bujarski S, Lau AS, Lee SS, Ray LA. Genetic and environmental predictors of alcohol use in Asian American young adults. *J Stud Alcohol Drugs* 2015;76:690-9.
  28. Wall TL, Luczak SE, Hiller-Sturmhöfel S. Biology, genetics, and environment: underlying factors influencing alcohol metabolism. *Alcohol Res* 2016;38:59-68.
  29. Eng MY, Luczak SE, Wall TL. ALDH2, ADH1B, and ADH1C genotypes in Asians: a literature review. *Alcohol Res Health* 2007;30: 22-7.
  30. Luczak SE, Pandika D, Shea SH, Eng MY, Liang T, Wall TL. ALDH2 and ADH1B interactions in retrospective reports of low-dose reactions and initial sensitivity to alcohol in Asian American college students. *Alcohol Clin Exp Res* 2011;35:1238-45.
  31. Wall TL. Genetic associations of alcohol and aldehyde dehydrogenase with alcohol dependence and their mechanisms of action. *Ther Drug Monit* 2005;27:700-3.
  32. Baik I, Cho NH, Kim SH, Han BG, Shin C. Genome-wide association studies identify genetic loci related to alcohol consumption in Korean men. *Am J Clin Nutr* 2011;93:809-16.
  33. Dickson PA, James MR, Heath AC, Montgomery GW, Martin NG, Whitfield JB, et al. Effects of variation at the ALDH2 locus on alcohol metabolism, sensitivity, consumption, and dependence in Europeans. *Alcohol Clin Exp Res* 2006;30:1093-100.
  34. Kim DJ, Choi IG, Park BL, Lee BC, Ham BJ, Yoon S, et al. Major genetic components underlying alcoholism in Korean population. *Hum Mol Genet* 2008;17:854-8.
  35. Hendershot CS, Neighbors C, George WH, McCarthy DM, Wall TL, Liang T, et al. ALDH2, ADH1B and alcohol expectancies: integrating genetic and learning perspectives. *Psychol Addict Behav* 2009;23:452-63.
  36. Duranceaux NC, Schuckit MA, Luczak SE, Eng MY, Carr LG, Wall TL. Ethnic differences in level of response to alcohol between Chinese Americans and Korean Americans. *J Stud Alcohol Drugs* 2008;69:227-34.
  37. Li H, Borinskaya S, Yoshimura K, Kal'ina N, Marusin A, Stepanov VA, et al. Refined geographic distribution of the oriental ALDH2\*504Lys (nee 487Lys) variant. *Ann Hum Genet* 2009;73(Pt 3):335-45.
  38. Yukawa Y, Muto M, Hori K, Nagayoshi H, Yokoyama A, Chiba T, et al. Combination of ADH1B\*2/ALDH2\*2 polymorphisms alters acetaldehyde-derived DNA damage in the blood of Japanese alcoholics. *Cancer Sci* 2012;103:1651-5.
  39. Paik IH, Lee C, Lee SJ, Lee CU, Kim JJ, Jun TY, et al. Genetic frequencies of aldehyde dehydrogenase 2 in Korea patients with alcohol dependence. *J Korean Neuropsychiatr Assoc* 1998;37:149-58.
  40. Paik K. The studies on the differences in personality trait by aldehyde dehydrogenase-I genotype variances in Korean male alcoholics. *J Korean Neuropsychiatr Assoc* 1995;34:15-26.
  41. Hashimoto M, Watanabe M, Uematsu Y, Hattori S, Miyai N, Utsumi M, et al. Relationships of alcohol dehydrogenase 1B (ADH1B) and aldehyde dehydrogenase 2 (ALDH2) genotypes with alcohol sensitivity, drinking behavior and problem drinking in Japanese older men. *Environ Health Prev Med* 2016;21:138-48.
  42. Mulligan MK, Ponomarev I, Hitzemann RJ, Belknap JK, Tabakoff B, Harris RA, et al. Toward understanding the genetics of alcohol drinking through transcriptome meta-analysis. *Proc Natl Acad Sci U S A* 2006;103:6368-73.
  43. Kim SW, Bae KY, Shin HY, Kim JM, Shin IS, Youn T, et al. The role of acetaldehyde in human psychomotor function: a double-blind placebo-controlled crossover study. *Biol Psychiatry* 2010;67:840-5.
  44. Lee B, Nam B, Seo JS, Jang S, Lee S, Yi SH. Elimination rate and pharmacokinetics of alcohol in Korean healthy male adults. *J Korean Neuropsychiatr Assoc* 2015;54:427-34.
  45. Kawano Y. Physio-pathological effects of alcohol on the cardiovascular system: its role in hypertension and cardiovascular disease. *Hypertens Res* 2010;33:181-91.
  46. Brooks PJ, Enoch MA, Goldman D, Li TK, Yokoyama A. The alcohol flushing response: an unrecognized risk factor for esophageal cancer from alcohol consumption. *PLoS Med* 2009;6:e50.
  47. Assanangkornchai S, Noi-pha K, Saunders JB, Ratanachaiyavong S. Aldehyde dehydrogenase 2 genotypes, alcohol flushing symptoms and drinking patterns in Thai men. *Psychiatry Res* 2003;118: 9-17.
  48. Crabb DW, Edenberg HJ, Bosron WF, Li TK. Genotypes for aldehyde dehydrogenase deficiency and alcohol sensitivity. The inactive ALDH2(2) allele is dominant. *J Clin Invest* 1989;83:314-6.
  49. Kim JM, Shin IS, Kim SW, Lee H, Yoon BH, Yoon JS. Relationships between Lifetime alcohol consumption and cognitive function in an elderly population: effect modification by level of education. *Korean J Psychopharmacol* 2003;14:239-43.
  50. Sin IS, Yoon JS, Kim H, Lee HY, Yoon BH, Lee H, et al. Effects of alcohol on neurocognitive function, psychomotor performance and subjective response in Koreans with different ALDH2 genotypes. *J Korean Soc Biol Psychiatry* 1999;6:176-88.
  51. Zhao Y, Wang C. Glu504Lys single nucleotide polymorphism of aldehyde dehydrogenase 2 gene and the risk of human diseases. *Biomed Res Int* 2015;2015:174050.
  52. Gu JY, Li LW. ALDH2 Glu504Lys polymorphism and susceptibility to coronary artery disease and myocardial infarction in East Asians: a meta-analysis. *Arch Med Res* 2014;45:76-83.
  53. Wang Q, Zhou S, Wang L, Lei M, Wang Y, Miao C, et al. ALDH2 rs671 Polymorphism and coronary heart disease risk among Asian populations: a meta-analysis and meta-regression. *DNA Cell Biol* 2013;32:393-9.
  54. Jia K, Wang H, Dong P. Aldehyde dehydrogenase 2 (ALDH2) Glu504Lys polymorphism is associated with hypertension risk in Asians: a meta-analysis. *Int J Clin Exp Med* 2015;8:10767-72.
  55. Rehm J, Shield KD, Roerecke M, Gmel G. Modelling the impact of alcohol consumption on cardiovascular disease mortality for comparative risk assessments: an overview. *BMC Public Health* 2016; 16:363.
  56. Chen L, Smith GD, Harbord RM, Lewis SJ. Alcohol intake and blood pressure: a systematic review implementing a Mendelian randomization approach. *PLoS Med* 2008;5:e52.



57. Ota M, Hisada A, Lu X, Nakashita C, Masuda S, Katoh T. Associations between aldehyde dehydrogenase 2 (ALDH2) genetic polymorphisms, drinking status, and hypertension risk in Japanese adult male workers: a case-control study. *Environ Health Prev Med* 2016;21:1-8.
58. Yin G, Naito M, Wakai K, Morita E, Kawai S, Hamajima N, et al. ALDH2 polymorphism is associated with fasting blood glucose through alcohol consumption in Japanese men. *Nagoya J Med Sci* 2016;78:183-93.
59. Jo SA, Kim EK, Park MH, Han C, Park HY, Jang Y, et al. A Glu487Lys polymorphism in the gene for mitochondrial aldehyde dehydrogenase 2 is associated with myocardial infarction in elderly Korean men. *Clin Chim Acta* 2007;382:43-7.
60. Eom SY, Zhang YW, Kim SH, Choe KH, Lee KY, Park JD, et al. Influence of NQO1, ALDH2, and CYP2E1 genetic polymorphisms, smoking, and alcohol drinking on the risk of lung cancer in Koreans. *Cancer Causes Control* 2009;20:137-45.
61. Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 2003;32:1-22.
62. Shin C, Kwack K, Cho NH, Kim SH, Baik I. Sex-specific differences in the association of a common aldehyde dehydrogenase 2 gene polymorphism and alcohol consumption with stroke risk in a Korean population: a prospective cohort study. *Nutr Res Pract* 2015;9:79-86.
63. Yun KE, Chang Y, Yun SC, Davey Smith G, Ryu S, Cho SI, et al. Alcohol and coronary artery calcification: an investigation using alcohol flushing as an instrumental variable. *Int J Epidemiol* 2017 Jan 10 [Epub ahead of print]. <https://doi.org/10.1093/ije/dyw237>.
64. Zuccolo L, Holmes MV. Commentary: Mendelian randomization-inspired causal inference in the absence of genetic data. *Int J Epidemiol* 2016 Dec 26 [Epub ahead of print]. <https://doi.org/10.1093/ije/dyw327>.
65. Alcohol Project Supporting Committee. Alcohol statistics white paper. Goyang: Alcohol Project Supporting Committee; 2010.
66. Oberbeck N, Langevin F, King G, de Wind N, Crossan GP, Patel KJ. Maternal aldehyde elimination during pregnancy preserves the fetal genome. *Mol Cell* 2014;55:807-17.
67. Yokoyama A, Kumagai Y, Yokoyama T, Omori T, Kato H, Igaki H, et al. Health risk appraisal models for mass screening for esophageal and pharyngeal cancer: an endoscopic follow-up study of cancer-free Japanese men. *Cancer Epidemiol Biomarkers Prev* 2009;18:651-5.
68. Shin HY, Shin IS, Yoon JS. ALDH2 genotype-associated differences in the acute effects of alcohol on P300, psychomotor performance, and subjective response in healthy young Korean men: a double-blind placebo-controlled crossover study. *Hum Psychopharmacol* 2006;21:159-66.
69. Jee YH, Lee SJ, Jung KJ, Jee SH. Alcohol intake and serum glucose levels from the perspective of a mendelian randomization design: the KCPS-II biobank. *PLoS One* 2016;11:e0162930.
70. Xu L, Jiang CQ, Cheng KK, Au Yeung SL, Zhang WS, Lam TH, et al. Alcohol use and gamma-glutamyltransferase using a Mendelian randomization design in the Guangzhou Biobank Cohort Study. *PLoS One* 2015;10:e0137790.
71. Au Yeung SL, Jiang C, Long M, Cheng KK, Liu B, Zhang W, et al. Evaluation of moderate alcohol use with QT Interval and heart rate using Mendelian randomization analysis among older Southern Chinese men in the Guangzhou Biobank Cohort Study. *Am J Epidemiol* 2015;182:320-7.
72. Au Yeung SL, Jiang C, Cheng KK, Cowling BJ, Liu B, Zhang W, et al. Moderate alcohol use and cardiovascular disease from Mendelian randomization. *PLoS One* 2013;8:e68054.
73. Au Yeung SL, Jiang CQ, Cheng KK, Liu B, Zhang WS, Lam TH, et al. Evaluation of moderate alcohol use and cognitive function among men using a Mendelian randomization design in the Guangzhou Biobank Cohort Study. *Am J Epidemiol* 2012;175:1021-8.
74. Tabara Y, Arai H, Hirao Y, Takahashi Y, Setoh K, Kawaguchi T, et al. The causal effects of alcohol on lipoprotein subfraction and triglyceride levels using a Mendelian randomization analysis: The Nagahama Study. *Atherosclerosis* 2017;257:22-8.
75. Tabara Y, Ueshima H, Takashima N, Hisamatsu T, Fujiyoshi A, Zaid M, et al. Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. *Atherosclerosis* 2016;254:242-8.